

IMPACT OF NURSE-ASSISTED CARDIAC REHABILITATION INTERVENTIONS ON HEALTH BEHAVIOR MODIFICATION OF PATIENTS WITH MYOCARDIAL INFARCTION

Sidra Bibi, Sarfraz Masih, Muhmmad Afzal

Received: 21 December, 2023 Revised: 29 January, 2024 Accepted: 05 February, 2024 Published: 16 February, 2024

ABSTRACT

Background: The rate of coronary thrombosis, especially MI, is increasing in Pakistan day by day and cardiologists are using interventional technology like angioplasty to treat such diseases. But this treatment is not enough for the patients to remain healthy for a long time and to get back to their daily life activities. Cardiac Rehabilitation is now becoming an utmost requirement for the cardiac patients due to their needs of health behaviors modifications to increase their expected year of life, which is fulfilled by the collaboration of a cardiologist, nurse, physiotherapist and nutritionist.

Study design: Quasi experimental (pre and post)

Setting: The Lahore School of Nursing, UOL.

Methodology: The study employed Paired sample T-tests and descriptive statistics to examine changes in variables before and after cardiac rehabilitation.

Results: There was a significant improvement in patients' overall health behavior pre- to post-CR, Results showed a significant positive correlation between post-CR behaviour and physical activity ($r = 0.566$; $P = 0.000$), food intake ($r = 0.316$; $P = 0.001$), and medication adherence ($r = 0.428$; $P = 0.000$).

Conclusion(s): pre interventions data were obtained from MI patients there were not any noteworthy practice in physical activity, nutrition and medication adherence. By the conclusion of the interventions, however, we had seen statistically significant changes in all three aspects of health behaviors.

Key words: MI, Health behavior modifications, Physical activity, dietary habits, medication adherence, nurse assisted cardiac rehabilitation.

INTRODUCTION

A Cardiac disease that includes ischemic heart disease, stroke, heart failure and myocardial infarction are the most common cause of estimated 13 million deaths worldwide in 2020. Myocardial infarction is the major component of cardiovascular disease (CVD) and ranks sixth in terms of health loss (after falls, low back pain, and chronic obstructive pulmonary disease). MI comes in second to IHD as the national top cause of mortality (Wilson, Cleghorn, Nghiem, & Blakely, 2023).

A quarter of all fatalities and one in every five deaths 20 years earlier occur due to coronary thrombosis disease (Mc Namara, Alzubaidi, Jackson, & practice, 2019). An estimated 580,000 American adults have a myocardial infarction (MI) each year, and 86% of these people recover from the condition. Patients having a history of MI are more likely than the

general population to experience another MI or another cardiovascular (CV) illness.

American Heart Association recommendations advise MI survivors to take evidence-based drugs and outpatient cardiac rehabilitation (CR) to lower their risk of further CV occurrences (Bush et al., 2020). According to AHA exercise-based cardiac rehabilitation improves exercise capacity and lowers mortality in individuals with acute myocardial infarction (AMI). Additionally, it may enhance quality of life, cardiovascular functional capacity, and psychological profile (Kirolos et al., 2019).

To overcome mortality, morbidity, and unexpected hospital admissions rate Nurse assisted CR interventions have shown effectiveness to control unhealthy health behaviors by increasing the exercise capacity, dietary behavior modification, smoking

cessation, psychological well-being and medication adherence. Nurse assisted Cardiac rehabilitation is currently considered mandatory in worldwide guidelines. Initiation of cardiac rehabilitation intervention after acute myocardial infarction will decrease 90% of repeated cardiac events and mortality (Ögmundsdóttir Michelsen et al., 2022) . In light of this context, our study's goal was to rank the CVD risk factor domains first, taking into account both the possible health benefits and health-related costs of preventative measures, in addition to the extent of the health loss(Wilson et al., 2023).

The risk factors of myocardial infarction (MI) reveal a multifaceted landscape with both modifiable and non-modifiable contributors. Established non-modifiable factors include age, gender (with males generally at higher risk), and family history of cardiovascular diseases. Modifiable risk factors, on the other hand, play a crucial role in prevention. Extensive evidence links cigarette smoking to a significantly increased risk of MI. Smoking cessation emerges as a pivotal intervention. Elevated blood pressure remains a key risk factor, emphasizing the importance of regular monitoring, lifestyle modifications, and pharmacological interventions. Elevated levels of LDL cholesterol and reduced levels of HDL cholesterol contribute to atherosclerosis and subsequent MI. Lipid-lowering therapies play a central role in risk reduction(Medicine, 2023)

The association between diabetes and MI is well-established. Glycemic control and lifestyle modifications are essential in managing this risk factor, amplifying the risk of MI Sedentary lifestyles contribute to cardiovascular risk. Regular physical activity is protective and integral in preventing MI. Diets high in saturated fats, trans fats, and sodium, along with low intake of fruits and vegetables, contribute to atherosclerosis and MI risk. A heart-healthy diet is crucial to MI prevention. Interventions targeting both non-modifiable and modifiable risk factors, along with patient education and community-wide initiatives, are essential for reducing the incidence of myocardial infarction(Espinosa-Salas & Gonzalez-Arias, 2023).

Nurse assisted cardiac rehabilitation interventions is the application of critical thinking in patient care with collaboration of a rehabilitation team to help patients to improve, maintain and regain

their health, achieving the highest standard of life, secondary prevention, behaviour modifications like involve in physical activity, healthy diet and tobacco cessation, stress management and medication adherence to enhance healthy year of life, reduce mortality and readmission rates (Gutenbrunner, Stievano, Nugraha, Stewart, & Catton, 2022).

MATERIAL AND METHODS:

The design used in this study was a quasi experimental pre and post. The present study was carried out at Bahria International Hospital Lahore.because this hospital has cardiac rehabilitation center. A non-probability convenient sampling technique were used to select a sample size on 3rd day of admission after myocardial infarction. Sample size of 105 cases was calculated with 95% confidence interval, 20% margin response rate was added to control drop off rate and the final sample was 111 with desire precision 0.05 ,health behaviours modification (physical activity) among myocardial infarction patients(Arjunan, D'Souza, & Health, 2021).In Nurse assisted cardiac rehabilitation interventions nurse works in a collaboration with a team of cardiologist, dietitian and physiotherapist. Nurse as a primary researcher coordinate with team for facilitating the cardiac rehabilitation of Myocardial infarction patients about the health behavior modifications like.

- Level of physical activity
- Diet chart approved by dietician and cardiologist
- Medication adherence to control blood pressure

The SPSS version 26 was used to analyse the data. Frequency, percentage, mean and standard deviation were used to measure the data. descriptive analysis and paired T-test were carried out whereby the mean differences between measurements were analysed for pre and post both the interventional findings.

RESULTS

Introductory paragraph

This chapter includes the results of demographic characteristics of study participant, overall results of health behavior modification and individual results of health behavior modification in physical activity, dietary habits and medication Adherence.

There were 88 males and 23 females in the control group, aged from 18 to 60 years, with an average age of (39 ± 8) years. There was no significant difference

in gender, age and other data between the two groups ($P < 0.05$).

Table1 Results of Demographic Profile of study participant

Demographic variable		Gender		Pre interventions			Post interventions		
		Female	Male	Mean	N	SD	Mean	N	SD
Age in years	18--26	1(0.9)	3(2.7)	0.75	4	0.500	0.75	4	0.500
	27--35	4(3.6)	12(10.8)	0.75	16	0.447	0.75	16	0.447
	36--45	6(5.4)	26(23.4)	0.81	32	0.397	0.81	32	0.397
	46--54	5(4.5)	21(18.9)	0.81	26	0.402	0.81	26	0.402
	55--60	7(6.3)	26(23.4)	0.79	33	0.415	0.79	33	0.415
Educational level	Middle	1(0.9)	1(0.9)	1.00	2	1.414	1.00	2	1.414
	Matric	2(1.8)	6(5.4)	3.00	8	1.069	3.00	8	1.069
	Graduation	6(5.4)	39(35.5)	2.44	45	1.216	2.44	45	1.216
	Master and above	14(12.6)	42(37.8)	2.75	56	1.083	2.75	56	1.083

According to the study shown in Table 1, there were more male patients with myocardial infarction (MI) admitted to Bahria International Hospital (BIH), Lahore (79.2% of n=88 patients with a standard deviation of 0.407 and 20.7% of female patients (n=23 patients with a standard deviation of 0.407). Male MI incidence was highest among admission ratios for those between the ages of 36 and 45. 35.5% (n=39) with SD 1.216 had graduated from high school, which was followed by female.

Patients who were admitted to the cardiac rehabilitation unit and those who had finished a four-month rehabilitation program were examined for this study. The following variables pertaining to alterations in health behavior specific to a given disease were measured. This measure has 21 items that span three domains: medication adherence (six items), dietary habits (fourteen items), and physical activity related to myocardial infarction (MI) (seven items).

Table2 overall results of health behavior modification

Overall HBM	Scoring	Pre interventions frequency (%)	Post interventions frequency (%)
Good health behavior modification	59-87	1(0.9%)	10(9%)
Average health behavior modification	31-56	75(67.5%)	101(91%)

Low health behavior modification	1-28	35(31.5%)	0(0%)
No health behavior modification	0	0(0%)	0(0%)

Health behavior modification in PA		Pre interventions frequency (%)	Post interventions frequency (%)
No Activity /Week	No HBM=0	0(0%)	0(0%)
Mild Activity /Week	Mild HBM =1-7	68(61.3%)	0(0%)
Moderate Activity /Week	Moderate HBM=8-14	41(36.8%)	21(18.9%)
Vigorous Activity /Week	High HBM=15-21	2(1.8%)	90(81.1%)
Health behavior modification in DH			
Taking diet never as per advised /week	No HBM=0	0(0%)	0(0%)
Taking diet occasionally as per advised/week	Mild HBM =1-15	97(87.4%)	1(0.9%)
Taking diet mostly as per advised /week	Moderate HBM=16-30	14(12.6%)	11(9.9%)
Taking diet always as per advised /week	High HBM=31-48	0(0%)	99(89%)

Health behavior modification in MA			
Taking medicines always as per advised /week =	No HBM=0	1(0.9%)	4(3.6%)
Taking medicines mostly as per advised/week	Mild HBM =1-6	1(0.9%)	98(88.3%)
Taking occasionally as per advised /week	Moderate HBM=7-12	35(31.5%)	7(6.3%)
Taking medicines never as per advised /week	Good HBM=13-18	74(66.7%)	2(1.8%)

The overall findings of this study on the use of nurse-assisted cardiac rehabilitation to modify the health behaviors of patients who have experienced a myocardial infarction are shown in Table 3. The pre- and post-date frequency percentage comparison demonstrates the changes in low behavior and

average behavior modification practices, with 35 (31.5%) becoming 0% and 75 (67.5%) becoming 101 (91%) respectively. The percentage of people who modified their behaviors for optimal health went from 0.9% to 9%.



Table 3 Results of health behavior modification

According to the analysis of Table 3, weekly intense physical activity results in significant changes in health-related behavior. Pre data showed that just 2 patients (1.8%) engaged in weekly intense physical activity, but post data showed that 90 patients (81.1%) did so. When myocardial infarction patients follow weekly dietary recommendations, their dietary habits improve significantly. Before the intervention, 0(0%) of patients followed the recommended serving sizes; shortly after the, 99(89%) of patients followed the recommendations. To control Blood pressure among

myocardial infarction patients who never took their medications as prescribed was 74 (66.7%) in the pre-data and significantly improved in the post-data with a decrease in the number of patients 2(1.8%). The Nurse-Assisted Cardiac Rehabilitation Intervention has resulted in notable improvements in health behavior modification among myocardial infarction patients, particularly in areas such as physical activity, dietary habits, and medication adherence. These findings suggest the intervention's effectiveness in fostering positive changes in patient behavior for better health outcomes.

Health behavior modification in PA		Pre interventions frequency (%)	Post interventions frequency (%)
No Activity /Week	No HBM=0	0(0%)	0(0%)
Mild Activity /Week	Mild HBM =1-7	68(61.3%)	0(0%)
Moderate Activity /Week	Moderate HBM=8-14	41(36.8%)	21(18.9%)
Vigorous Activity /Week	High HBM=15-21	2(1.8%)	90(81.1%)
Health behavior modification in DH			
Taking diet never as per advised /week	No HBM=0	0(0%)	0(0%)
Taking diet occasionally as per advised/week	Mild HBM =1-15	97(87.4%)	1(0.9%)
Taking diet mostly as per advised /week	Moderate HBM=16-30	14(12.6%)	11(9.9%)
Taking diet always as per advised /week	High HBM=31-48	0(0%)	99(89%)
Health behavior modification in MA			
Taking medicines always as per advised /week =	No HBM=0	1(0.9%)	4(3.6%)
Taking medicines mostly as per advised/week	Mild HBM =1-6	1(0.9%)	98(88.3%)
Taking occasionally as per advised /week	Moderate HBM=7-12	35(31.5%)	7(6.3%)
Taking medicines never as per advised /week	Good HBM=13-18	74(66.7%)	2(1.8%)

Table 4 Results of Physical Activity Domain of Health Behavior Modification

Physical Activity days /week						
		Mean interventions (SD)	Pre Mean interventions (SD)	Post Mean interventions (SD)	T Value	Sig
Q1	Vigorous PA	1.06(0.856)		1.93 (0.723)	0.566	0.000
Q2	Moderate PA	1.05(0.562)		2.31(0.585)	0.317	0.001
Q3	Mild PA	0.90(0.617)		2.29(0.624)	0.217	0.222
physical Activity time /week						
Q1	Vigorous PA	1.08(0.605)		2.53(0.615)	0.152	0.112
Q2	Moderate PA	0.95(0.743)		2.11(0.652)	0.386	0.000
Q3	Mild PA	1.03(0.744)		2.32(0.618)	0.357	0.000
Q4	No PA	1.30(0.838)		2.59(0.610)	0.314	0.001

Subsequently in table 04 presented data of MIPDM (myocardial infarction, physical activity, dietary habits and medication adherence) shows distribution of pre and post analysis related to physical activity as mild to vigorous in a days per week. Table shows that vigorous physical activity days/week is increased in the examined group, the mean vigorous physical activity was 1.06 (0.856) before nurse assisted cardiac rehabilitation intervention and becomes 1.93 (0.723) after rehabilitation. The PA max increased significantly after NACR ($p = 0.000$) (correlation coefficient=0.566). Some comparative

characteristics for this phase of rehabilitation are presented above in table 4.

The physical activity time/week, ranging from no physical activity to intense exercise and arranged as pre- to post-data analysis. Table shown that following nurse-assisted cardiac rehabilitation interventions, frequency of moderate and mild physical activity per week were increased as Pre interventions data shows mean were 0.95(0.625) and after interventions 2.11 (0.652) with (correlation coefficient=0.386) ($p = 0.000$). Some comparative characteristics for this phase of rehabilitation are presented above in Table 4.

Table05 Results of Dietary servings/week of health behavior modification

Servings/week		Mean Pre interventions (SD)	Mean Post interventions (SD)	T Value	Sig
Q1	Seeds, nuts and dry fruits	1.27(0.904)	2.57(0.582)	0.27	0.003
Q2	servings of vegetables	1.21(0.728)	2.60(0.561)	0.337	0.000
Q3	servings of fruits	0.89(0.608)	2.37(0.617)	0.277	0.003
Q4	Servings of whole grains	1.00(0.661)	2.46(0.584)	0.165	0.084
Q5	Fat free dairy product	0.95(0.519)	2.36(0.600)	0.180	0.059
Q6	Meat, fish and poultry	0.69(0.629)	2.20(0.536)	0.316	0.001
Q7	servings of fats and oil	0.85(0.621)	2.40(0.561)	0.033	0.731
Q8	servings of sugar-sweetened	0.82(0.635)	2.31(0.553)	0.159	0.096

In dietary servings table 05 the paired sample *t*-test showed that there was no significant follow-up of DASH diet in myocardial infarction patient before interventions, but nurse assisted cardiac Rehabilitation interventions led to a noteworthy increased in taking of advised servings of food. As it is revealed in the pre-controlled group the

recommended servings of fruits per week with mean were 0.89(0.608) and increased to 2.37(0.617) (*P* 0.277) with significance level of 0.003 in post-group. The mean of weekly servings of vegetable before intervention were 1.21(0.728) increased after intervention to 2.60 (0.617) with significance value of 0.000.

Table06 Results of Dietary servings/day of health behavior modification

Servings/day		Mean Pre (SD)	Mean Post (SD)	T Value	Sig
Q1	servings of whole grains	0.86(0.653)	2.40(0.561)	0.98	0.306
Q2	servings of vegetables	0.85(0.575)	2.32(0.587)	0.359	0.000
Q3	servings of fruits	0.77(0.583)	2.41(0.562)	0.142	0.136
Q4	servings of fat free product	0.80(0.585)	2.30(0.597)	0.040	0.676
Q5	servings of meat, fish poultry	0.87(0.634)	2.41(0.610)	0.395	0.000
Q6	servings of sugar - sweetened	0.95(0.706)	2.35(0.627)	0.385	0.000

One of the most often recommended dietary plans for lowering blood pressure is the diet to stop hypertension in myocardial infarction patients. A dietary strategy to lower blood pressure may work by

raising plasma nitrite levels to improve the ability of vascular endothelium to upregulate NO (nitric oxide). In actuality, because nuts, dry fruits and fish are high in L-arginine, they can raise NO levels. It

can also enhance vascular reactivity due to its strong antioxidant content. Our goal in this study was to

help MI patients adhere to the DASH diet for a duration of six months.

Table 7 Results of Medication Adherence of Health Behavior Modification

Medication/week		Mean Pre (SD)	Mean Post (SD)	T Value	Sig
Q1	Forget to take medicine	2.20(0.772)	0.92(0.764)	0.428	0.000
Q2	Have problem to remember	2.27(0.82)	0.86(0.720)	0.417	0.000
Q3	How many days stop to take medicine	2.33(0.705)	0.59(0.623)	-0.041	0.666
Q4	Feel worse to take medicine	2.18(1.055)	0.64(0.671)	-0.126	0.189
Q5	medication of your own free choice	2.32(0.765)	0.65(0.656)	-0.43	0.657
Q6	Feel more normal on medication	2.15(1.020)	0.81(0.707)	-0.010	0.918

The goal of cardiac rehabilitation is to assist patients that have recently suffered a cardiac event, such as myocardial infarction (MI), by developing a routine in a safe, monitored environment. The 3rd component of health behavior modifications in cardiac rehabilitation centers is medication adherence that is demarcated as the amount to which

individuals take their medicines as directed through doctors to improve and maintain cardiac health. Medication non-adherence was common, especially after MI as pre data revealed that patients forget to take medicine mean were 2.20 (0.772) after nurse assisted interventions improved to 0.92 (0.764) with (R=0.428) and (P= 0.000).

DISCUSSION

In the discussion chapter of this thesis, we delve into a comprehensive analysis and interpretation of the research findings, aiming to provide valuable insights into the implications of our study. By synthesizing the results with existing literature, we illuminate the significance of our contributions to the field and address the research questions, thereby offering a nuanced understanding of the broader context surrounding our investigation, includes discussion on demographic picture of study participant, overall results of health behavior modification and individual results of physical activity domain, dietary habits domain and medication adherence.

The findings of this groundbreaking study which focused on patient education for cardiac rehabilitation (CR) of myocardial infarction patients, demonstrated notably higher patient behavior scores following an education intervention. The goal of Nurse assisted cardiac rehabilitation is to

assist patients that have recently suffered a cardiac event, such as myocardial infarction (MI) in physical activity, dietary habits and medication intake. Similar results were shown by other studies that used health behaviors as their outcomes (de Melo Ghisi et al., 2020; Mansilla-Chacon et al., 2021; Su, Yu, & Paguio, 2020). These studies support the use of nursing education interventions in conjunction with cardiac rehabilitation (CR) as a standard of care for myocardial infarction patients, while also highlighting the significance of CR as a cornerstone of care. The design of this study and its findings demonstrated that, despite variations in program settings and treatments, applying study results to specific programs may be a challenging task. Another study shows that total of eighteen post-MI patients (mean age: 60.5 years, range 37–73 years) took part in the study. Participants were predominantly male (n = 13, 72 %)(Coull & Pugh, 2021).in contrast to this study where there were 88

males and 23 females in the control group, aged from 18 to 60 years, with an average age of (39 ± 8) years. The study's results further supported the link between socioeconomic position and health, as evidenced by the fact that participants with higher educational status had greater effects than their counterparts and that educational treatments improved the patients' ability to modify their health-related behaviors. It's also critical to emphasize that the individuals in sample were male and had greater levels of education, which may have contributed to their participation in these programs and referrals. Education is a crucial aspect of CR; thus, it should be taken into account along with sex and age (Su et al., 2020).

This study shows MI incidence in Male between aged 36-45 and 55-60 years of age were high. Male with graduation and master level of education were also high amongst admission ratios. In this study education intervention were translated into both Urdu and English. From the standpoint of the patient, participants were eager to learn and pleased with the instruction given. The majority expressed satisfaction and confidence in the information they had been given.

A systematic assessment of secondary preventive educational interventions, including a thorough overview and quantitative synthesis of the efficacy of disease-related information and CHD risk reduction behaviors in adults with MI diagnoses. The findings of this research show that different forms and intensities of nurse assisted educational interventions for secondary prevention can effectively improve healthy behaviors and disease awareness at the <6 and 6-12-month follow-up. The most remarkable findings concerned the short-term (less than six months) enhancement of healthy eating practices, with patients with CHD three times more likely to follow a healthy diet following the delivery of structured educational interventions. There was a notable shift in the amount of physical activity and medication adherence, and there was a chance that these would be maintained (Shi et al., 2023).

The overall findings of this study on the use of nurse-assisted cardiac rehabilitation to modify the health behaviors of patients who have experienced a myocardial infarction on 4-months follow-up of nursing interventions shows that overall improvement in health behavior modification related

to physical activity, diet and medication adherence. The pre- and post-date frequency percentage comparison demonstrates the percentage of people who modified their behaviors for optimal health went from 0.9% to 9%.

Increased local and systemic blood flow, shear stress, and nitric oxide release from moderate and more vigorous exercise all contribute to the enhancement of chronic flow-mediated dilatation. patients who have had a myocardial infarction have much reduced blood flow. However, it appears that increased physical activity might at least significantly mitigate this drop of blood flow. Physical activity leads to vessel stiffness is somehow concerned in older age above 75 years (Tršan, Košuta, Fras, & Jug, 2021). The results of paired sample-t-test of this study shows that vigorous physical activity everyday/week is increased in the intervention group, the mean of vigorous physical activity was 1.06 (0.856) before nurse assisted rehabilitation intervention and 1.93 (0.723) after rehabilitation. The PA max increased significantly after NACR ($p = 0.000$) (correlation coefficient=0.566). Similarly moderate physical activity per week were increased as Pre interventions data shows mean were 0.95(0.625) and after interventions 2.11 (0.652) with (correlation coefficient=0.386) ($p = 0.000$).

A randomized controlled trial (RCT) shows the beneficial effects of a 6-month behavioral nutrition intervention that was started in the clinic and included phone and mail follow-ups that emphasized the DASH diet on systolic blood pressure and vascular function. DASH-4-Teen's participants achieved a substantial increase in DASH dietary adherence from pre- to post-intervention comparison to adolescents receiving RC by increasing their intake of fruits, low-fat dairy foods, and associated nutrients and decreasing their intake of total and saturated fat and salt. In comparison to RC patients, who saw a more moderate change in DASH adherence and systolic blood pressure at the end of therapy, mean (SD) baseline intake of these same dietary components (daily fruit servings: 1.3 [1.4] versus 0.7 [0.7], $P=0.01$, and percent of calories from fat: 31.8 [5.8] versus 34.4 [5.6], $P=0.01$). baseline daily fruit servings (mean [SD]: 1.2 [1.3] versus 0.6 [0.7], $P=0.04$) and percent of calories from fat (32.1 [6.0] versus 34.8 [4.5], $P=0.04$) (Couch et al., 2021). DASH diet the best had a reduced risk of heart failure

than those who adhered the least Cronbach's alpha of the scale was found to be 0.72 (Ibsen et al., 2022).

The paired sample-t-test showed that there was no significant follow-up of DASH diet in myocardial infarction patient at the baseline, The mean of weekly servings of vegetable before intervention were 1.21(0.728) increased after intervention to 2.60 (0.617) with significance value of (0.000). As the daily servings of meat, fish and poultry in myocardial infarction according to DASH diet plan were poor as mean of pre data 0.87(0.634) in post data mean was 2.41(0.610) with ($P = 0.000$) and (coefficient of correlation = 0.395), it means that cardiac rehabilitation interventions play a significant role in the dietary management of myocardial infarction patients.

Medication Adherence Scale (MMAS-6) was assessed using Pearson's correlation, which showed a correlation of 0.6851 ($p < 0.01$). The Cronbach's α of the scale is 0.61, supporting its internal consistency reliability. A p-value of less than 0.05 was considered statistically significant (Zheng et al., 2020). In my study Medication non-adherence was common, especially after MI as pre data revealed that patients forget to take medicine mean were 2.20 (0.772) after nurse assisted interventions improved to 0.92 (0.764) with ($R=0.428$) and ($P= 0.000$).

CONCLUSION:

At the baseline, data were obtained from myocardial infarction (MI) patients on admission there were not any noteworthy practice in physical activity, nutrition and medication adherence and majority of the patients had no excellent health behavior. By the conclusion of the interventions, however, we had seen statistically significant changes in all three aspects of health behavior by IPAQ, DASH diet, and MMAS-6. The advantages of an education interventions have been confirmed in this first-ever site research in Lahore, Pakistan, which focuses on patient education for physical activity, diet and medication adherence.

REFERENCES:

- Ács, P., Veress, R., Rocha, P., Dóczy, T., Raposa, B. L., Baumann, P., . . . Makai, A. J. B. P. H. (2021). Criterion validity and reliability of the International Physical Activity Questionnaire–Hungarian short form against the RM42 acceleromter. *21*(1), 1-10.
- Arjunan, P., D'Souza, M. S. J. C. E., & Health, G. (2021). Efficacy of nurse-led cardiac rehabilitation on health care behaviours in adults with chronic heart failure: An experimental design. *12*, 100859.
- Bush, M., Kucharska-Newton, A., Simpson Jr, R. J., Fang, G., Stürmer, T., Brookhart, M. A. J. J. o. c. r., & prevention. (2020). Effect of initiating cardiac rehabilitation after myocardial infarction on subsequent hospitalization in older adults. *40*(2), 87.
- Couch, S. C., Saelens, B. E., Khoury, P. R., Dart, K. B., Hinn, K., Mitsnefes, M. M., . . . Urbina, E. M. J. H. (2021). Dietary approaches to stop hypertension dietary intervention improves blood pressure and vascular health in youth with elevated blood pressure. *77*(1), 241-251.
- Coull, A., & Pugh, G. J. B. c. d. (2021). Maintaining physical activity following myocardial infarction: a qualitative study. *21*, 1-9.
- de Melo Ghisi, G. L., Rouleau, F., Ross, M.-K., Dufour-Doiron, M., Belliveau, S. L., Brideau, J.-R., . . . Oh, P. J. C. o. (2020). Effectiveness of an education intervention among cardiac rehabilitation patients in Canada: a multi-site study. *2*(4), 214-221.
- Espinosa-Salas, S., & Gonzalez-Arias, M. (2023). Behavior Modification for Lifestyle Improvement. In *StatPearls [Internet]*: StatPearls Publishing.
- Gutenbrunner, C., Stievano, A., Nugraha, B., Stewart, D., & Catton, H. J. I. n. r. (2022). Nursing—a core element of rehabilitation. *69*(1), 13-19.
- Ibsen, D. B., Levitan, E. B., Åkesson, A., Gigante, B., & Wolk, A. J. E. J. o. P. C. (2022). The DASH diet is associated with a lower risk of heart failure: a cohort study. *29*(7), 1114-1123.
- Kirolos, I., Yakoub, D., Pendola, F., Picado, O., Kirolos, A., Levine, Y. C., . . . Khouzam, R.

- N. J. A. o. T. M. (2019). Cardiac physiology in post myocardial infarction patients: the effect of cardiac rehabilitation programs—a systematic review and update meta-analysis. *7*(17).
- Mansilla-Chacon, M., Gomez-Urquiza, J. L., Martos-Cabrera, M. B., Albendin-Garcia, L., Romero-Bejar, J. L., Canadas-De La Fuente, G. A., . . . Disease. (2021). Effects of supervised cardiac rehabilitation programmes on quality of life among myocardial infarction patients: a systematic review and meta-analysis. *8*(12), 166.
- Mc Namara, K., Alzubaidi, H., Jackson, J. K. J. I. p. r., & practice. (2019). Cardiovascular disease as a leading cause of death: how are pharmacists getting involved? , *8*, 1.
- Medicine, G. C. R. C. J. N. E. J. o. (2023). Global effect of modifiable risk factors on cardiovascular disease and mortality. *389*(14), 1273-1285.
- Ögmundsdóttir Michelsen, H., Sjölin, I., Bäck, M., Gonzalez Garcia, M., Olsson, A., Sandberg, C., . . . Leósdóttir, M. J. J. o. M. I. R. (2022). Effect of a lifestyle-focused web-based application on risk factor management in patients who have had a myocardial infarction: randomized controlled trial. *24*(3), e25224.
- Shi, W., Ghisi, G. L., Zhang, L., Hyun, K., Pakosh, M., & Gallagher, R. J. J. o. C. N. (2023). Systematic review, meta-analysis and meta-regression to determine the effects of patient education on health behaviour change in adults diagnosed with coronary heart disease. *32*(15-16), 5300-5327.
- Su, J. J., Yu, D. S. F., & Paguio, J. T. J. J. o. A. N. (2020). Effect of eHealth cardiac rehabilitation on health outcomes of coronary heart disease patients: A systematic review and meta-analysis. *76*(3), 754-772.
- Tršan, J., Košuta, D., Fras, Z., & Jug, B. J. F. i. P. (2021). Vascular function in patients after myocardial infarction: the importance of physical activity. *12*, 2367.
- Wilson, N., Cleghorn, C., Nghiem, N., & Blakely, T. J. P. H. M. (2023). Prioritization of intervention domains to prevent cardiovascular disease: a country-level case study using global burden of disease and local data. *21*(1), 1.
- Zheng, F., Ding, S., Lai, L., Liu, X., Duan, Y., Shi, S., & Zhong, Z. J. F. i. P. (2020). Relationship between medication literacy and medication adherence in inpatients with coronary heart disease in Changsha, China. *10*, 1537.